



PHYSICOCHEMICAL CHARACTERISTICS AND PLANKTON ASSEMBLAGES OF SATHIYAR RIVER IN MADURAI DISTRICT

Vanitha.M¹ and Joseph Thatheyus.A²

¹ Department of Zoology, SBK College, Aruppukottai, Tamil nadu, India.

² PG & Research Department of Zoology, The American College, Madurai, Tamil nadu, India.

Abstract:

Water is an important natural resource on earth. It is necessary for all living organisms, ecological systems, human health, food production and economic development. Water can be obtained from a number of sources, among which are streams, lakes, rivers, ponds, rain, springs, and wells. The ensuring of good quality drinking water is a basic factor in guaranteeing public health, protection of environment and sustainable development. Urbanization has direct impact on water bodies as human settlement takes place around the vicinity of water bodies causing encroachment near water bodies. People obtain their consumable water from surface and ground water. However, both surface and ground water sources could become polluted by biological and chemical contaminants arising from point and non-point sources. Surface water is polluted with high levels of nutrients (Nitrogen and Phosphorus) which cause eutrophication, leading to the growth of other microorganisms and causes contamination in surface water. Other contamination sources of surface water are household wastes, sewage, industrial effluent, synthetic detergents, agrochemicals, and oil spillage. The raising of pollutants in the river results in health hazards to aquatic life directly

or indirectly by altering physico-chemical parameters. Since these sources are open, they are highly susceptible to flood and birds, animals and human contamination. The quality of water is affected by an increase in anthropogenic activities and any pollution either physical or chemical causes changes in the quality of the receiving water body. Hence, the present study has been designed to evaluate the physico-chemical parameters and related plankton assemblage in Sathiyar river. The river under study lies geographically between latitude 10 0 4'0" and longitude 10 0 4'0". It originates from sirumalai hills (a group of hills on the Eastern Ghats) situated 25km away from Dindigul town. For irrigation purpose, a dam has been constructed across this river namely Sathiyar Dam in Palamedu. The overflow of water in the Sathiyar Dam during rainy season empties into Vaigai River. In this preliminary study, six sites along the Sathiyar river were randomly selected. The physico-chemical parameters such as colour, odour, turbidity, TDS, EC, pH, alkalinity and salinity were analyzed. In addition to this, the plankton species were also collected in these six sites. The correlation between these two data were statistically analyzed and interpreted to find the influence of physico-chemical parameters on plankton assemblage in the Sathiyar River.

Keywords: Sathiyar river, Water quality, Physico-chemical parameters, Plankton assemblage, and Pollution

Introduction

River water is not only an indispensable source for irrigation but also plays a vital role for drinking water supply for most of the urban municipalities. Water from rivers is pumped at specific sites and after treatment at municipal water treatment plants supplied as domestic potable water supply (Shah and Singh 2016). Water pollution is an acute problem in all the major rivers and dams. Water is known to contain a large numbers of chemical element, and the interactions of both the physical and chemical properties of water play a major role in composition, distribution and abundance of aquatic organisms. Water is extensively used for irrigation, industrial development, hydro electrical generations, fisheries, human life survival and for domesticated animals. Hence physico-chemical analysis is of prime importance to assess the quality of water for its best usage and also to know the pollution load on receiving water bodies. The maintenance of healthy aquatic ecosystem is dependent on the physicochemical properties and biological diversity. Among these, plankton constitute the

foundation of the food web in aquatic ecosystems and represent one of the most direct and profound responses to pollution entering river. Phytoplanktons are also the indicators of water pollution. Several researchers informed on a number of algal species as water quality indicator (Naik et al., 2005; Nandan and Aher, 2005; Zargar and Ghosh, 2006). In a study on Kadra reservoir of Karnataka in 2006, Zargar and Ghosh reported on several algal forms belonging to Chlorophyceae, Cyanophyceae, and Bacillariophyceae as water pollution indicators. Nandan and Aher (2005) concluded that the algal genera, *Oscillatoria*, *Scenedesmus* and *Navicula* are the species found in organically polluted waters. Hence, the present study has been designed to evaluate the water quality and related plankton assemblage in Sathiyar River.

Materials and method

The water samples for hydrobiological studies, phytoplankton and zooplankton were collected at six selected sampling sites of the Sathiyar River. The distance between two adjacent sites is approximately 2kms. Sampling was carried out weekly random water sampling method. The samples were taken at the same time each day in the upper reach of the river in downstream direction. Sampling was carried out for three months of monsoon period from June to August 2015. In each sampling sites, the sample water was collected with a 100 ml plastic bottle. The water samples for phytoplankton were preserved with 1% Lugol's iodine solution. Qualitative and quantitative analyses were carried out in the laboratory using the U termol method under an inverted microscope (Utermohl, 1958). The phytoplankton composition was based on the identification of the specific and infra-specific taxa. Zooplankton sampling at each station was collected with standard plankton net of 1 m length, 30 cm mouth diameter, and mesh size of 1m fitted with a flow meter. Samples were preserved in a 4% buffered formalin/river water solution. Zooplankton species were identified to the lowest taxonomic unit possible, and taxon abundance (per cubic meter) was counted under a microscope (1 ml sub sample). These samples were taken with a Stempel-pipette from the entire sample (250 ml).

The water samples from the six selected sites (collected sites are sathiyar dam, Erampatti, Kovilpatti, Ayyur, Kuravankulam and Vandiyur respectively) were periodically collected and the physico-chemical parameters such as Turbidity, TDS, EC, pH, Total hardness Ca, Mg, Fe, and Nitrites, Nitrates, Fluorides and Chlorides were estimated (APHA, 1995). Simultaneously

the phytoplankton and zooplankton were also estimated. The estimated values were statistically analyzed.

Results and Discussion

Safe water supply is recognized at the highest priority task in environmental protection throughout the globe (Adewuyi et al., 2014). For the present study, the water samples of all six selected sites (collected sites are sathiyar dam, Erampatti, Kovilpatti, Ayyur, Kuravankulam and Vandiyur respectively) were periodically collected and the physico-chemical parameters such as Turbidity, TDS, EC, pH, Total hardness as CaCO_3 , Ca, Mg, Fe, and Nitrite as NO_2 , Nitrate as NO_3 , Fluoride and chlorides were estimated. Simultaneously the phytoplankton and zooplankton were also estimated. The estimated values were statistically analyzed and tabulated in Table-1 and 2. According to the table values, all the physico-chemical parameters observed for sites I, II, III and IV were more or less similar when compared to the sites V and VI. In the present study, the mean values of pH of Sathiyar river varied between 7.4 to 8.3. These values indicated that pH of drinking water source was slightly alkaline in nature. In the current study, slight alkaline nature of Sathiyar river is in accordance with the previous reports of (Mullai et al., 2013) while evaluating pH of Uppanar river of Tamilnadu. In sites V and VI, the physico-chemical parameters such as Turbidity, TDS, EC, Total hardness as CaCO_3 , Ca, Mg and chlorides were one fold elevated than that of the other sites. The other parameters such as nitrites and nitrates were meagerly increased in the sites V and VI when compared to other sample sites of the river. Even though considerable increase occurred in both sites V and VI, the values of physico-chemical parameters observed in all samples sites were occurred within the acceptable and the permissible limits. The elevation of parameters Turbidity, TDS, EC, Total hardness as CaCO_3 , Ca, Mg and chlorides in the sites V and VI may be due to the mixing of non-point source of sewage and polluted water into these two regions of Sathiyar River. This finding has some coincidence with the TDS, TH, turbidity and Cl values found in Gudbahri river and important indicators of pollution (Weldemariam 2013).

The results for plankton numbers were inversely proportionate to the results of physico-chemical parameters. The fluctuations in the total number of plankton in all four sites such as I, II, III and IV were low when compared to the sites V and VI. The sites V and VI showed the decreasing trends in the number of plankton and also disappearance of plankton

communities such as *Cladophora*, *Coelosphaerium*, *Calanoids nauplii* and *Brachionus*. This may be due to discharge of pollutants in these sites. The decomposition of the organic materials by micro-organisms in the aquatic ecosystem leads to the lowering of the level of dissolved oxygen, which in turn inhibits the growth or cause the death of the aquatic habitats and the planktonic population depends on hydrological situation as well as water quality parameters. Water quality in an aquatic ecosystem is determined by many physical and chemical factors (Sargaonkar & Deshpande 2003). The poor diversity of plankton may be caused by poor light penetration. Algal analysis thus showed that water quality of the pond has reached at threshold level and therefore, it needs some corrective measures to maintain the water chemistry of the river (Nair et al., 2015). Basic water management principles have to be designed to maintain good water quality and reduce incidence of diseases. It is important to develop a plan of action to be taken when water quality parameters are outside the desirable range and in stressful concentrations. Hence monitoring regularly and recording data is important as it will aid in anticipation of needed action.

REFERENCES

- [1] Aedwuyi, G. O., Etchie, A.T. and T.O. Etchie (2014). Health risk assessment of exposure to metals in a nigerian water supply. *Human and Ecological Risk Assessment: An international Journal* 20, 29-44.
- [2] APHA (1995). *Standards methods for the examination of water and wastewater*. 19th edition., American Public Health Association, New York.
- [3] Asha M.S. Nair, J.K. Reshma, Anu Mathew and J.A Aswathy Ashok (2015). Effect of Water Quality on Phytoplankton Abundance in Selected Ponds of Edumangad Block Panchayat, Kerala. *Emer Life Sci Res.*, 1(2): 35-40.
- [4] Mullai P., Yogeswari, M.K.,Oommen, B.,Sarvanakumar,K., and K. Kathiresan, (2013). Monitoring of water quality parameters in Uppanar river of Cuddalore district, Tamilnadu state, India. *Journal of Water Sustainability*, 3(4), 179-192.
- [5] Naik, U.G., Bhosale, S.H., Rathod J.L., and U. G. Bhat (2005). Diversity of phytoplankton groups in the river Kali, west coast of India. *Proc. of the State Level UGC Sponsored Seminar on Biodiversity and its Conservation, Haveri, 28-29th July*, pp. 192-196.

- [6] Nandan., S.N. and A. H. Aher (2005). Algal community used for assessment of water quality of Haranbaree dam and Mosam river of Maharashtra. *J. Environ. Biol.*, 26: 223-227
- [7] Sargaonkar A. and V. Deshpande (2003). Development of an Overall Index of Pollution for surface water based on a general classification scheme in the Indian context. *Environ. Monit. Assess.*, 89 (1), 43–67
- [8] Shah A. B. and R.P. Singh (2016). Monitoring of Hazardous Inorganic Pollutants and Heavy Metals in Potable Water at the Source of Supply and Consumers end of a Tropical Urban Municipality. *Int. J. Environ. Res.*, 10(1):149-158.
- [9] Utermohl, H (1958). Zur Vervollkommnung der qualitativen Phytoplanktonethodik. *Mit Int Theor Angew Limnol*, 9: 1-38.
- [10] Weldemariam M.M., (2013). Physico-chemical analysis of GudBahri River Water of Wuro, Eastern Tigray, Ethiopia. *International Journal of Scientific and Research Publications*, 3(11). ISSN 2250-3153.
- [11] Zargar S. and T.K. Ghosh (2006). Influence of cooling water discharges from Kaiga nuclear power plant on selected indices applied to plankton population of Kadra reservoir. *J. Environ. Biol.*, 27: 191-198.