



SCIREA Journal of Clinical Medicine

ISSN: 2706-8870

<http://www.scirea.org/journal/CM>

August 8, 2023

Volume 8, Issue 4, August 2023

<https://doi.org/10.54647/cm321137>

Predicting of Poor Outcomes for Patients HIV-associated Community-Acquired Pneumonia: Who are at risk and how reduce mortality in such patients?

Alizamin Sadigov ^{1,*}, Rauf Baylarov ¹, Malahat Sultanova ¹

¹ Pulmonary and Critical Care Medicine Department, Therapeutic and Education Clinic, Medical University of Azerbaijan, Baku. Azerbaijan

*Correspondence: alizaminsadigov@gmail.com

Abstract

Community-acquired pneumonia (CAP) in individuals with HIV results from multiple risk factors, particularly immune defects and a CD4 count decrease, especially when below < 100 cells/mm³ continue to be a risk for pneumonia due to routine bacterial pathogens. Assessment of risk factors which are responsible for poor prognosis in patients with CAP associated with HIV infection is important and interesting for further evaluation of possible their modification and through that reduction mortality rate and improved the patient outcomes. We aimed to investigate the risk factors predicting poor outcomes in patients with CAP developed in individuals with HIV infection. We conducted a retrospective cohort study involved patients with diagnosed CAP who were admitted to the pulmonary and critical care unit of medical university teaching hospital, Baku city, Azerbaijan, from January of 2018 to December of 2022. One hundred ninety three adult patients (>18 years old) with CAP were enrolled to the study. The number of patients with CAP associated with HIV infection was 59 (30.9% of all CAP patients) and 16 of them we have identified accidentally in our hospital. In

our study recurrent pneumonia episodes ,especially in patients with $CD4 < 500$ cells/mm³ was higher and the severity of immunodeficiency has predicted the frequency and severity of recurrent pneumonia episodes(9.5[2.8-18/7]; $p < 0.01$) and the $Cd4 < 100$ cells/mm³ count was major risk factor predicting ICU admission and bacteremia with severe sepsis/septic shock (20.9[10.4-30.8]; $p < 0.001$).We have found five major and independent risk factors predicting in-hospital mortality in our study:severe malnutrition; $CD4 < 100$ cells/mm³;radiographic progression of the disease; $P/F < 250$;and bacteremia with severe sepsis/septic shock .We identified factors associated with an increased risk of ICU admission and fatal outcomes which could help identify patients who might benefit from anti-pneumococcal vaccination ,antiretroviral treatment ,adequate nutrition and ventilatory support as well as determine prognosis.Our findings should be validated by studies with larger samples of patients.

Keywords: HIV infection, community acquired pneumonia, risk factors , ICU admission, poor outcomes, mortality

Introduction

Community-acquired pneumonia CAP in individuals with HIV results from multiple risk factors, particularly immune defects ACD4 count decrease, especially when below < 100 cells\mm³ continues to be a risk factor for pneumonia due to routine bacterial pathogens.

Other immune defects include quantitative and qualitative B-cell abnormalities that result in impaired pathogen specific antibody production, abnormalities in neutrophil function and members, and abnormalities in alveolar macrophage function. Lack of anti-retroviral therapy CART or intermittent use of ART increases the risk of pneumonia, likely due to uncontrolled HIV viremia.

Additional risk factors that contribute to the continued risk for bacterial pneumonia in individuals with HIV include chronic viral hepatitis, tobacco, alcohol, injection drug use and prescribed opioid use, particularly higher doses and opioids with immunosuppressive properties. Chronic obstructive pulmonary disease [COPD], malignancy, renal replacement therapy, and congestive heart failure (CHF) are risk factors for pneumonia, particularly in the population of older adults with HIV.

In individuals with HIV the incidence of bacteremia accompanying pneumonia is greater than in individuals without HIV, especially when infection is due to *S.pneumoniae* with the introduction of ART and pneumococcal conjugate vaccines for both the general pediatric population and individuals living with HIV, this disparity in incidence rates of bacteremia between people without HIV has narrowed but has not been eliminated. In one recent study of invasive pneumococcal disease (IPD), which includes bacteremia, IPD was more common in people with HIV who had CD4 counts <500 cells/mm³, but even those with counts >500 cells/mm³, had a higher incidence than in the general population .

Assessment of risk factor which may predict poor outcomes for patients with CAP in individuals with HIV is important for further evaluation of possibility they modification and through that reduction mortality rate and improved patient outcomes we aimed to investigate the risk factors for CAP in patients with HIV infection and poor outcomes in such patients.

Methods

We conducted a retrospective cohort study involving patients with diagnosed CAP who were admitted to the pulmonary and critical care medicine unit at Medical University hospital, Baku City, Azerbaijan, between January of 2018 and December of 2022. One hundred ninety three adults (>18 years) patients with CAP were enrolled to the study

Pneumonia was diagnosed as CAP when it occurs before 48h of hospital admission 59 of 193 patients with CAP pneumonia was developed in individuals with HIV infection ,and 16 of them the HIV infection was diagnosed in our hospital .The criteria of diagnosis are new persistent pulmonary infiltrates appearing on chest radiographs and a least two of following :

(1) Fever of >38°C, (2) leukocytosis of >12.000/mm³ or <4.000/mm³ and (3) purulent sputum. In case of clinically suspected pneumonia sputum and blood culture (in case of fever>38°C) was sent for microbiology and positive quantitative culture (cut of point > 10~6CFV/ml was assessed. To analyze the predisposing factors and patients outcomes the following variables were evaluated: age, gender, comorbidities, the presence of HIV infection at the time of hospitalization, length of hospital stay, ICU admission rate, mechanical ventilation rate, development of complications and in hospital mortality.

All comorbidities were defined as previously described. Diagnosis of malnutrition was assessed based on the presence of one or more of the following criteria :body mass index

less than $<21\text{kg}/\text{m}^2$;serum albumin concentrations $<3.5\text{g}/\text{L}$. Septic shock was defined as bacteremia and hypotension requiring use of vasopressors with presence of poly-organ dysfunction. The severe pneumonia required ICU admission was defined with septic shock required use of vasoactive drugs and need to mechanical ventilation. Multi drug resistant (MDR) pathogens was defined as isolation of bacterial strain non-susceptible to at one agent in three one more antimicrobial categories.

The study end points were the assessment risk factors for ICU admission and in hospital mortality. To identify risk factors associated with ICU admission and in hospital mortality were performed univariable and multivariable analysis to detect significant differences between groups we used the chi square test or fishers exact test for categorical variables and the two tailed test or Mam-Whitney test for continuous variables when appropriate continuous variables were reported as medians with interquartile ranges (IQR) :members and percentages were reported for categorical variables. Base line variables (recorded at pneumonia onset and hospital admission) were analyzed in the univariable analysis to identify ICU admission and in-hospital mortality. Statistical significance was established of <0.5 .

Result

In total, 193 patients with CAP were analyzed. The number of patients with CAP in individuals with HIV infection was 59 (30.9% of all CAP incidence) and 16 of them HIV infection was identified in our hospital .

Table 1 describes baseline characteristics of our study population.

Table 1. Baseline characteristics of patients with CAP with-without HIV infection

	Over all n=193%	CAP without HIV n=134%	CAP with HIV n=59%	P value
Males	117(60.0)	81(60.4)	36(61.0)	0.41
Females	76(40.0)	53(39.5)	23(39.0)	0.42
Age, median years	51 {32-76}	58 {42-76}	44 {32-56}	<0.01
Prior antibiotic therapy (last90 days)	60(31.0)	21(15.7)	39(66.1)	<0.01
Chronic heart	45(23.3)	32(23.1)	13(22.0)	0.28

failure				
Renal replacement therapy	36(18.6)	11(8.2)	25(42.4)	<0.01
COPD	38(19.6)	25(18.6)	13(22.0)	0.11
Bronchiectasis	29(15.0)	8(6.0)	21(35.0)	<0.5
Malnutrition	44(22.8)	10(5.2)	34(57.6)	<0.03
Diabetes	52(26.9)	39(29.1)	13(22.0)	0.9
Pleural effusion	59(30.5)	28(20.8)	31(52.5)	<0.01
Bilateral pneumonia	49(25.3)	21(15.6)	28(47.4)	<0.04
Unilateral pneumonia	144(74.7)	113(84.3)	31(52.6)	<0.02
Lobar consolidation	82(42.5)	69(51.4)	13(22.0)	<0.02
Focal segmental consolidation	111(57.5)	65(48.5)	46(77.9)	<0.04
Cavitation	42(21.7)	19(14.1)	23(39.0)	<0.05
Bacteremia	51(26.4)	26(19.4)	25(42.4)	<0.01
CD4<500cells/mm ³	48(24.9)	9(6.7)	39(66.1)	<0.001
Recurrent pneumonia	42(21.8)	8(5.8)	34(57.6)	<0.01
Lack of pneumococcal vaccination	144(74.6)	92(68.6)	52(88.9)	<0.01
MDR pathogens	40(20.7)	19(14.2)	21(35.5)	<0.01
Severe sepsis/septic shock	55(28.5)	23(17.1)	22(39.0)	<0.01
ICV admission	60(31.0)	26(19.4)	34(57.6)	<0.02
In hospital mortality	39(20.2)	18(13.4)	21(35.5)	<0.03

There was significant differences between age of patients recording HIV infection. The patients with CAP in individuals with HIV was younger compared to patients with CAP without HIV (44[32-56]) vs 58[42-76]; $p<0.01$). Prior use of antibiotics (last 90 days) was

common in patients with CAP in individuals with HIV (15.7% vs 66.1%; $p<0.01$) and commonly it was related to the presence of persistent high fever in individuals with HIV before development of CAP. Hemodialysis also was common event in patients with CAP in individuals with HIV ($p<0.1$) and chronic use of renal replacement therapy (RRT) in such patients was related to catheter -driven blood stream infection (bacteremia) and development of MDR-pathogen associated pneumonia ($p<0.1$). Bronchiectasis as comorbidity was common in HIV-associated CAP patients ($p<0.5$) and it was also predict the prevalence of MDR-Pathogen associated pneumonia in such patients. Malnutrition was common demonstrated in HIV-associated CAP patients ($p<0.03$), and more severe malnutrition was associated with increased incidence of ICU admission of patients .

Unilateral and lobar consolidation frequently was found in patients with CAP ($p<0.02$), whereas bilateral and focal, segmental consolidation commonly was presented in patients with HIV -associated CAP (<0.04). Cavitation as radiographic feature of pneumonia has commonly found in HIV-associated pneumonia related to *S.aureus and P.aeruginosa* infections($p<0.01$)

We have found several risk factors which were predictable for developing bacteremia in our study and bacteremia as sign of invasive pneumonia was found frequently in HIV -associated CAP patients (42.4% vs 19.4%; $p<0.01$). Lymphopenia (CD4<500 cells/mm³) was common finding in HIV- associated CAP patients ($p<0.001$) and predict the severity of CAP.

More severe lymphopenia was associated with higher incidence of ICU admission and in hospital mortality of patients ($p<0.01$).

Lack of pneumococcal vaccination leads to clinically and radiological more severe pneumonia and the number of patients vaccinated against pneumococcal disease was significantly lesser in patients with HIV -associated CAP(88.1% vs 68.6%; $p<0.01$)

Pathogens were identified in 65 of 193 patients (54.4%), streptococcus pneumonia and *Hemophiles species* were most common frequently identified causes of CAP ,and were the same in individuals with and without HIV (Tabl 2).

Tabl 2.Pathogens isolated among 106 patients with culture-positive CAP

Etiologies	Over all n=106(%)	CAP without HIV n=64%	CAP with HIV n=42(%)	P value

<i>Streptococcus Pneumoniae</i>	3030(28.3)	18(21.1)	12(28.6)	0.44
<i>Haemophilus influenza</i>	18(11.3)	8(12.5)	4(9.5)	0.19
<i>Mycoplasma Pneumoniae</i>	10(11.3)	9(14.1)	1(2.3)	<0.05
<i>Chlamydia pneumoniae</i>	8	8(12.5)	--	NA
<i>Klebsiella pneumoniae</i>	2(1.9)	1(1.6)	1(2.3)	NA
<i>MSSA</i>	4(3.9)	2(3.3)	2(4.5)	NA
<i>Eserichia coli MDR pathogens</i>				
<i>Pseudomonas Aeruginosa</i>	18(16.9)	8(6.7)	10(23.8)	<0.01
<i>MRSA</i>	13(19.2)	4(6.3)	9(11.4)	<0.01
<i>Esherichia coli</i>	4(3.7)	1(1.6)	3(7.1)	NA
<i>Klebsiella pneumoniae</i>	5(4.7)	2(3.3)	NA	NA

However, atypical bacterial pathogens such as *Mycoplasma pneumoniae* and *Chladmidophilia pneumoniae* were common in patients with CAP without HIV infection ($p < 0.5$). The frequency of MDR-pathogens such as *Pseudomonas aeruginosa* and *Staphylococcus Aureus* caused CAP was higher in patients with HIV infection and as usual in such patients were several risk factors for MDR-pathogens associated CAP including poorly controlled HIV related to lack of ART and frequently use of antibiotics before admission to hospital for CAP. In our study there was played role previously structural lung disease such as bronchiectasis in patients with CAP developed in individuals with HIV ($p < 0.5$). As known exacerbations of bronchiectasis required frequently use of antibiotics which is associated with development of MDR pathogens. In our study gravity

of MDR pathogens was higher compared to pathogens with CAP without HIV (21(35.5) vs 19(14.2); $p < 0.1$)

Bacteremia and severe sepsis/septic shock were common among patients with CAP in individuals HIV (tab.3)

Table3. Multivariate analysis of in bacteremia and severe sepsis/septic shock in patients with CAP associated with HIV infection

Variables	OR	95%CL	P value
CD<100cells/mm ³	20.9	10.4-30.8	<0.001
Lack of ART	7.2	2.4-13.6	<0.01
Lack of pneumococcal vaccination	3.4	1.2-7.6	<0.01
MDR pathogens	2.7	1.0-5.7	<0.02
Severe malnutrition	3.1	2.9-5.6	<0.01
Hemodialysis	2.6	0.9-5.2	<0.01

Low CD4 level was associated with an increased risk of bacteremia and severe sepsis with ICU admission and it was strongly independent risk factor for severe pneumonia in patients with ICU admission (20.9[10.4-30.8]); $p < 0.001$)

Lack of ART in HIV infected patients as usual is associated with poorly controlled infection and viremia. However, it is also was associated with increased risk of severe sepsis and bacteremia (7.2[2.4-13.6]; $p < 0.01$)

In our study bacteremia and severe sepsis together with severe respiratory failure ($PaO_2/FiO_2 < 250$) were associated with ICU admission of patients ($p < 0.02$).

Mortality was higher in patients with CAP associated with HIV infection, and multivariable analysis showed that severe malnutrition (serum albumin level < 2.5 g/l, 31[0.8-5.8]; $p < 0.05$), CD4<100 cells/mm³ (10.4[2.8-19.6]; $p < 0.01$); radiographic progression of disease (3.2[0.99-6.7]; $p < 0.01$); severe respiratory failure ($PaO_2/FiO_2 < 250$; 3.5[1.1-6.9]; $p < 0.1$); and bacteremia and severe sepsis/septic shock (8.4[2.8-18.9; $p < 0.01$) were associated an increased risk of in hospital mortality of patients (Table 4).

Table 4. Multivariate analysis of in hospital mortality in CAP patients associated with HIV infection

Variables	OR	95%CL	P value
Severe malnutrition	3.1	0.8-5.8	<0.05
CD4<100 cells/mm ³	10.4	2.8-19.6	<0.01
Radiographic progression of disease	3.2	0.94-6.7	<0.01
PaO ₂ /FiO ₂ <250	3.5	1.1-6.9	<0.01
Bacteremia and severe sepsis	8.4	2.8-18.9	<0.01

Combination of risk factors significantly an increased death rate among HIV infection associated with CAP patients (Figure 1).




Figure 1. Combination of risk factors and mortality rate among patients with CAP associated HIV infection.

The presence of all five risk factors in one patients was associated with highly incidence of in hospital death and decreased the number of risk factors predicted mortality was associated with an increased survival of patients and single risk factor was associated with less mortality late

Discussion

Bacterial pneumonia is a common cause of HIV-associated morbidity. Recurrent pneumonia, considered two or more episodes within all year period, is an AIDS defining condition. The incidence of bacterial pneumonia in individuals with HIV has decreased progressively with the advent of combination antiretroviral therapy (ART [17-19]).

In our study recurrent pneumonia episodes, especially in patients with $CD4 < 500 \text{ cells/mm}^3$ was higher and the severity of immunodeficiency has predicted the frequency recurrent pneumonia (9.5[2.8-18.7]; $p < 0.01$). Our study has not shown the decreased incidence of CAP in individuals with HIV because lack of ART was common in our patients and it was associated with an increased incidence of viremia and poorly controlled HIV infection which was predicted higher incidence of CAP (7.2[2.4-13.6]; $p < 0.01$)

However despite ART, bacterial pneumonia remains more common in people with HIV than in those who do not have HIV. Bacterial pneumonia may be the first manifestation of underlying infection and can occur at any stage of HIV disease and at any CD4 count. In our study, the $CD4 < 500 \text{ cells/mm}^3$ level was major risk factor for bacterial pneumonia, however the $CD4 < 100 \text{ cells/mm}^3$ level was major risk factor predicting ICU admission and bacteremia with sepsis /septic shock (20.9[10.4-30.8]; $p < 0.001$)

The clinical and radiographic presentation in individuals with HIV, particularly in those with higher CD4 count and HIV viral suppression is similar to that in individuals without HIV. In contrast in our study there were significant differences between clinical and radiographic features of bacterial pneumonia with and without HIV infection. Lack of auscultative signs, such as egophony and fine crackles and bilateral focal segmental consolidation was common in HIV associated CAP patients. Compared to CAP patients in HIV associated CAP patients pleural effusion and cavitation were common and was caused by increased incidence of MDR *S.aureus* and *P.aeruginosa* infections. In patients with CAP unilateral and lobar consolidation was common findings ($p < 0.02$)

Although some studies suggest that bacterial pneumonia is associated with increased mortality in individuals with HIV, others do not. Independent predictors of increased mortality in a prospective multicenter study of individuals with HIV community acquired bacterial pneumonia were $CD4 < 100 \text{ cells/mm}^3$, radiographic progression of disease and presence of shock. In our study the presence of HIV infection in CAP patients with associated with an increased risk of ICU admission was associated with an increased risk of

in hospital mortality (2.5[0.94-5.6]); $p < 0.02$). We have found five major and independent risk factor predicting in hospital mortality in such patients; severe malnutrition; $CD4 < 100 \text{ cells/mm}^3$; radiographic progression of disease $P/F < 250$; and bacteremia with severe sepsis/sepsis shock. In our study the presence of HIV-infection in CAP patients with associated with an increased risk of ICU-admission and ICU admission was associated with an increased risk of in-hospital mortality (2.5[0.94-5.6]; $p < 0.02$). We have found five major and independent risk factor predicting in-hospital mortality in such patients: severe malnutrition; $CD4 < 100 \text{ cells/mm}^3$; radiographic progression of disease ; $P/F < 250$; and bacteremia with severe sepsis/septic shock. Comparison to reference analysis data we have found two additional risk factors which highly associated with an increased in hospital mortality in HIV-associated CAP patients (3.1[0.8-50.8]; $p < 0.5$ and 3.5 [1.1-6.9]; $p < 0.1$; respectively).

The present study has several limitations. First, it is a study carried out in a single center. Second, some variables and co-founders may not have been taken into account given the difficulty of including them in a retrospective study. Third, for microbiological identification we used sputum, blood, pleural effusion and in some intubated patients we use endobronchial aspirate which are characterized with low accuracy for definition of bacterial pathogens in mechanical ventilated patients the use of bronchoalveolar lavage (BAL) fluid and bronchial brush? from affected area of lung would associated with an increased incidence of causative pathogen identification. Fourth, the sample size was somewhat small, so external validity might have been compromised and the results might not be fully generalizable for some of the findings. This highlights need to for multicenter studies that address the particular aspects of CAP patients in individuals with HIV- infection.

In conclusion, we found a several risk factors in our sample, related to development of bacterial pneumonia in individuals with HIV-infection. We found a high burden of comorbidities mostly related to chronic kidney failure and severe malnutrition as well as considerably high in-hospital mortality in such patients. We identified factors associated with increased risk of ICU admission and fatal outcomes which could help identify patients who might benefit from anti-pneumococcal vaccination, adequate nutrition and antiretroviral therapy as well as determine prognosis. These findings should be validated by studies with larger samples of patients.

Reference

- [1] Rurshtivk RE, Glassroth J, Jordan MC, et al. Bacterial pneumonia in persons infected with the human immunodeficiency virus. Pulmonary complications of HIV infection study group. *N Engl J Med*. 1995;333(13): 845-851. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7651475>,
- [2] Sogaard OS, Lohse N, Gerstoft J. Hospitalization of pneumonia among individuals with and without HIV infection, 1995-2007: a Danish population based nationwide cohort study. *Clin Infect Dis*. 2008; 47(10): 1345-1353. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18834317>
- [3] Aston SJ, Ho A, Jary H, et al. Aetiology and risk factors for mortality in an adult community acquired pneumonia cohort in Malawi. *Ann Respir Crit Care Med*. 2019. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30625278>
- [4] Jambo KC, Banda DH, Kan Kwatira, AM, et al. Small alveolar macrophages are infected preferentially by HIV and exhibit impaired phagocytosis function. *Mucosal Immunol*. 2024;7(5):1116-1126. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24472847>
- [5] Charles TP, Shelito J. Human immunodeficiency virus infection and host defence in the lungs. *Semin Respir Crit Care Med* 2016;37(2):
- [6] Gordin FM, Roediger MP, Girard P, et al. Pneumonia in HIV-infected persons: increased risk with cigarette smoking and treatment interruption. *Am J Respir Crit Care Med*. 2008; 178 (6): 630-636.
- [7] Lamas CC, Coelho LE, Grinsztejn BJ, Veloso VG. Community-acquired lower respiratory tract infections in HIV-infected patients on antiretroviral therapy: predictors in a contemporary cohort. *Infection*. 2017;45(6):801-809.
- [8] Edelman ES, Gordon KS, Crothers K, et al. Association of Prescribed Opioids With Increased Risk of Community-Acquired Pneumonia Among Patients With and Without HIV. *JAMA Intern Med*. 2019;179(3):297-304.
- [9] Burack JH, Hohn JA, Saint-Maurice D, Jacobson MA. Microbiology of community-acquired bacterial pneumonia in persons with and at risk for human immunodeficiency virus type 1 infection. Implications for rational empiric antibiotic therapy. *Arch Intern Med*. 1994; 154(22): 2589-2596.
- [10] Müller RF, Foley NM, Kessel D, Jeffrey AA. Community-acquired lobar pneumonia in patients with HIV infection and AIDS. *Thorax*. 1994; 49(4): 367-368

- [11] Mundy LM, Auwaerter PG, Oldach D, et al. Community-acquired pneumonia: impact of immune status. *Am J Respir Crit Care Med.* 1995 ; 152(4 Pt1): 1409-1315.
- [12] Rimland D, Navin TR, Lennox JL, et al. Prospective study of etiologic agents of community-acquired pneumonia in patients with HIV infection. *AIDS.* 2002; 16 (1): 85-95.
- [13] Afessa B, Green B. Bacterial pneumonia in hospitalized patients with HIV infection: the Pulmonary Complications, ICU Support, and Prognostic Factors of Hospitalized Patients with HIV (PIP) Study. *Chest* 2000; 117-1022
- [14] Attia EF, McGinns KA, Feemster LC, et al. Association of COPD with risk for pulmonary infections requiring hospitalization in HIV infected veterans. *J Acquir Immune Defic Syndrom* 2015; 70(3): 280-288.
- [15] Muccini C, Galli L, Lepri AC, et al. Incidence, timing, and determinants of bacterial pneumonia among HIV- infected patients: data from the ICONA Foundation cohort. *J Acquir Immune Defic Syndr.* 20B; 63 (3): 339-345.
- [16] Bordon JM, Fernandez-Bortan R, Wiemken TL, et al. Bacteremic pneumococcal pneumonia: clinical outcomes and preliminary results of inflammatory response. *Infection.* 2015; 43(6): 729-738.
- [17] Zolopa A, Anderson J, Powderly W et al. Early antiretroviral therapy reduces AIDS progression/death in individuals with acute opportunistic infections: a multicenter randomized strategy trial. *PLoS One.* 2009; 4(5): e55 75.
- [18] O'Connor J, Vjecha MJ, Philips AN, at al. Effect of immediate initiation of antiretroviral therapy on risk of severe bacterial infections in HIV-positive people with CD4 cell counts of more than 500 cells per μ L: secondary outcome results from a randomised controlled trial. *Lancet HIV* 2017; 4(3): e105- e112.
- [19] Sullivan JH, Moore RD, Keruly JC, Chaisson RE. Effect of antiretroviral therapy on the incidence of bacterial pneumonia in patients with advanced HIV infection. *Am J Respir Crit Care Med* 2000; 162(1): 64-67.
- [20] Cilloniz C, Torres A, Manzardo C, et al. Community-acquired pneumococcal pneumonia in virologically suppressed HIV-infected adult patients: a matched case control study. *Chest.* 2017; 152 (2): 295-303.